arXiv:0807.4829v1 [math.GR] 30 Jul 2008

WHEN ARE DUAL CAYLEY AUTOMATON SEMIGROUPS FINITE?

VICTOR MALTCEV

Mathematical Institute, University of St Andrews
St Andrews, Fife KY16 9SS, Scotland, `victor.maltcev@gmail.com`

Abstract. In this note we prove that, for a finite semigroup S, the dual Cayley automaton semigroup $\mathbf{C}^*(S)$ is finite if and only if S is $\mathcal{H}$-trivial and has no non-trivial right zero subsemigroups.

Keywords: automaton semigroup, Cayley automaton.

1. Introduction & Main Theorem

In a recent paper [6], Silva & Steinberg for any finite semigroup S define the *Cayley automaton* $\mathcal{C}(S)$ associated with S: its state set is S; the alphabet, on which the states act, is S; and when $\mathcal{C}(S)$ is in state s and reads symbol x, it moves to the state sx and outputs the symbol sx. In other words, this is the automaton obtained from the Cayley graph of S, specifying the output symbol on the arc, going from s and labeled by x, to be sx.

As to every automaton, to the automaton $\mathcal{C}(S)$ we can associate its automaton semigroup $\mathbf{C}(S)$, i.e. the transformation semigroup on infinite sequences S^∞, generated by states $s \in S$ viewed as the correspondent transformations of S^∞. In [6] Silva & Steinberg prove the following

Theorem 1.1. *Let G be a finite non-trivial group. Then $\mathbf{C}(G)$ is a free semigroup of rank $|G|$.*

The author and Alan Cain after reading [6] decided to study semigroups $\mathbf{C}(S)$ in more details. In [1] we characterized when $\mathbf{C}(S)$ is free, commutative, trivial, a left or a right zero semigroup. But the most interesting result we obtain in [1] is:

Theorem 1.2. *Let S be a finite semigroup. Then $\mathbf{C}(S)$ is finite if and only if S is $\mathcal{H}$-trivial.*

The sufficiency of Theorem 1.2 was proved in a recent work [4]. While the author and Alan Cain where preparing [1], we did not know about the work [4]. The sufficiency of Theorem 1.2 is the harder bit of this result. We provided in [1] two different proofs of sufficiency of Theorem 1.2: using the method of actions on sequences and using the method of wreath recursions. These two approaches and the method from [4] are all different in their realizations but bear the same spirit: the aim was to find an ultimate constant N such that any product of states from $\mathcal{C}(S)$ of length N can be reduced to a shorter word.

Now, if in the Cayley graph of a finite semigroup S, instead of the output symbol sx we put the symbol xs, we arrive at the *dual Cayley automaton* $\mathcal{C}^*(S)$: the state and alphabet sets being S and when $\mathcal{C}^*(S)$ is in state s and reads symbol x, it moves to the state sx and outputs the symbol xs.

Denote by $\mathbf{C}^*(S)$ the automaton semigroup generated by the automaton $\mathcal{C}^*(S)$. The main result of the note is

Theorem 1.3. *Let S be a finite semigroup. Then $\mathbf{C}^*(S)$ is finite if and only if S is $\mathcal{H}$-trivial and does not contain non-trivial right zero subsemigroups.*

We prove Theorem 1.3 in Section 3. The way we do it differs from the way of proving Theorem 1.2: there will be no need to deal with 'long' words and trying to reduce their lengths. The author hopes to exhibit the power of the method of wreath recursions.

Before we start the proof, in the next section we give all the auxiliary notation and lemmas we will need.

2. Auxiliary Lemmas

In order to distinguish the states and symbols in $\mathcal{C}^*(S)$, we will write $\overline{s}$ to denote the state correspondent to $s \in S$. If $S = \{s_1, \dots, s_n\}$, it is convenient to realize $\overline{s}$ via the *wreath recursion* (see [5]):

$$\overline{s} = \rho_s(\overline{ss_1}, \dots, \overline{ss_n}),$$

where $\rho_s : S \to S$, given by $x \mapsto xs$, corresponds to the action of $\overline{s}$ on the symbols S, and $\overline{ss_i}$ is the state where $\overline{s}$ moves after reading the symbol s_i.

If $x \in S$ and $\alpha \in \mathbf{C}^*(S)$ then by $q(\alpha, x)$ we will denote the state, to which α moves after reading $x \in S$. The transition function of α on S we will denote by $\tau(\alpha)$. So, $\tau(\alpha) : S \to S$ and for all $x \in S$, $x\tau(\alpha)$ is the symbol which outputs $\mathcal{C}^*(S)$, reading x in the state α. By definition, for all $a_1, \dots, a_k, x \in S$, we have

$$q(\overline{a_1} \cdots \overline{a_k}, x) = \overline{a_1 x} \cdot \overline{a_2 x a_1} \cdots \overline{a_k x a_1 \cdots a_{k-1}}. \tag{2.1}$$

Also the corresponding transition function $\tau(\overline{a_1} \cdots \overline{a_k})$ is $\rho_{a_1} \cdots \rho_{a_k} = \rho_{a_1 \cdots a_k}$.

Lemma 2.1. *Let S be a finite semigroup. Then for all $s, t \in S$, $\overline{s} = \overline{t}$ in $\mathbf{C}^*(S)$ if and only if $\rho_s = \rho_t$.*

Proof. Let $s, t \in S$. Then $\overline{s} = \overline{t}$ if and only if $\rho_s = \rho_t$ and $\overline{sx} = \overline{tx}$ for all $x \in S$. Recursing the latter, we obtain that $\overline{s} = \overline{t}$ if and only if $\rho_s = \rho_t$ and $\rho_{sx} = \rho_{tx}$ for all $x \in S$. It remains to notice that if $\rho_s = \rho_t$, then $\rho_{sx} = \rho_{tx}$ for all $x \in S$. □

In the following three lemmas we collect some information about $\mathbf{C}^*(S)$ for specific types of S.

Lemma 2.2. *Let G be a non-trivial finite group. Then $\mathbf{C}^*(G)$ is infinite.*

Proof. Take any non-identity element $a \in G$. Then $H = \langle a \rangle$ is a non-trivial commutative group. Obviously the restrictions of the action of the state $\overline{h}$, $h \in H$, to H^∞, is the same as the action of $\overline{h}$ in $\mathbf{C}^*(H)$. Notice that $\mathcal{C}(H) = \mathcal{C}^*(H)$ and so, by Theorem 1.1, $\overline{H}$ is a free system. Thus $\mathbf{C}^*(G)$ is infinite. □

Lemma 2.3. *Let L be a finite left zero semigroup and S be a finite semigroup. Then $\mathbf{C}^*(S \times L) = \mathbf{C}^*(S)$.*

Proof. Let $s \in S$ and $r, t \in L$. Then it follows from Lemma 2.1 that $\overline{(s, r)} = \overline{(s, t)}$ in $\mathbf{C}^*(S \times L)$. Hence $\mathbf{C}^*(S \times L)$ coincides with $T = \{\overline{(s, r_0)} : s \in S\}$ for any fixed $r_0 \in L$. It is now easy to check that $\overline{(s, r_0)} \mapsto \overline{s}$ gives rise to an isomorphism from T onto $\mathbf{C}^*(S)$. □

Lemma 2.4. *Let R be a finite right zero semigroup with $|R| > 1$. Then $\mathbf{C}^*(R)$ is a free semigroup of rank $|R|$.*

Proof. Let $R = \{q_1, \dots, q_n\}$ with $n > 1$. A direct calculation shows that $\mathcal{C}^*(R)$ is the automaton $\mathcal{A}$ of the type:



We will prove now that the automaton semigroup generated by the automaton $\mathcal{A}$ is free of rank n. If $\mathcal{A}$ has just read a symbol i, then it has just entered state q_i and its *next* output symbol will be i. Thus the action of q_i is to send a sequence α to $i\alpha$: the sequence α is shifted right by one symbol and the symbol i is inserted at the start. So if $w = q_{i_1} \cdots q_{i_m}$ with $i_j \in \{1, \dots, n\}$, then

$$1^\infty \cdot w = i_m i_{m-1} \cdots i_2 i_1 1^\infty \quad \text{and} \quad 2^\infty \cdot w = i_m i_{m-1} \cdots i_2 i_1 2^\infty.$$

The common prefix of $1^\infty \cdot w$ and $2^\infty \cdot w$ thus determines w and so the automaton semigroup generated by $\mathcal{A}$ is free with the basis R. ☐

The following lemma is standard to prove, we include it for completeness.

Lemma 2.5. *Let S be a finite semigroup and let $a, b \in S$. If all a, b and ab belong to the same $\mathcal{D}$-class of S, then $ab \in R_a \cap L_b$.*

Proof. By [2, Theorem 2.4] we have that $L_a R_b$ is contained in the same $\mathcal{D}$-class of S. Take any element $c \in R_a \cap L_b$. Then $c^2 \mathcal{D} c$. Let $\mu : S \to \mathcal{T}_X$ be any faithful representation of S into $\mathcal{T}_X$, the transformation semigroup on X, for some *finite* X. Obviously H_c is a group $\mathcal{H}$-class in S if and only if $H_{\mu(c)}$ is a group $\mathcal{H}$-class in $\mathcal{T}_X$. We have that $\mu(c)$ and $\mu(c)^2 = \mu(c^2)$ are $\mathcal{D}$-equivalent in $\mathcal{T}_X$. This happen only if $\mu(c) \mathcal{H} \mu(c)^2$ in $\mathcal{T}_X$. So that $H_{\mu(c)}$ is a group and so H_c is a group in S. Therefore $c \mathcal{H} c^2$ and so $L_a \cap R_b$ has an idempotent. By Clifford-Miller Theorem it now follows that $ab \in R_a \cap L_b$. ☐

3. PROOF OF THEOREM 1.3

Proof of Theorem 1.3. First notice that the condition that S does not contain non-trivial right zero subsemigroups is equivalent to the condition that there are no two distinct idempotents $e, f \in S$ such that $e \mathcal{R} f$.

($\Rightarrow$). Suppose that $\mathbf{C}^*(S)$ is finite.

S is $\mathcal{H}$-trivial.

Take any $\mathcal{H}$-class H in S. With the seek of a contradiction, suppose that $|H| > 1$. Let $T = \{t \in S : Ht \subseteq H\}$. Then for every $t \in T$, by [2, Lemma 2.21], the mapping $\gamma_t : h \mapsto ht$, $h \in H$, is a bijection of H onto itself. The set of all these bijections forms the so-called *Schützenberger group* $\Gamma(H)$ of H. By [2, Theorem 2.22] we have $|\Gamma(H)| = |H|$. Take arbitrary $t_1, \dots, t_k \in T$. Then, by (2.1), for all $x \in H$ we have:

$$q(\overline{t_1} \cdots \overline{t_k}, x) = \overline{t_1 x} \cdot \overline{t_2 x t_1} \cdots \overline{t_k x t_1 \cdots t_{k-1}}.$$

We also have $\tau(\overline{t_1} \cdots \overline{t_k}) = \tau(\overline{t_1 \cdots t_k})$.

Take now $\gamma_{t_1}, \dots, \gamma_{t_k}, \gamma_x \in \mathbf{C}^*(\Gamma(H))$. Then

$$q(\overline{\gamma_{t_1}} \cdots \overline{\gamma_{t_k}}, \gamma_x) = \overline{\gamma_{t_1} \gamma_x} \cdot \overline{\gamma_{t_2} \gamma_x \gamma_{t_1}} \cdots \overline{\gamma_{t_k} \gamma_x \gamma_{t_1} \cdots \gamma_{t_{k-1}}}$$

and $\tau(\overline{\gamma_{t_1}}\cdots\overline{\gamma_{t_k}}) = \tau(\overline{\gamma_{t_1}\cdots\gamma_{t_k}})$. Notice also that $\gamma_x\gamma_y = \gamma_{xy}$ for $x, y \in T$.

Take $t \in T$ and consider the restriction of $\overline{t}$ to H^∞. From the very definition of $\Gamma(H)$, it follows that the mapping $\overline{t}\restriction_{H^\infty} \mapsto \overline{\gamma_t}$ gives rise to a well-defined homomorphism from $\langle \overline{t}\restriction_{H^\infty} : t \in T\rangle$ onto $\mathbf{C}^*(\Gamma(H))$. By Lemma 2.2, it means that $\langle \overline{T}\rangle$ is infinite. Thus $\mathbf{C}^*(S)$ is infinite, a contradiction.

There are no distinct idempotents $e, f \in S$ such that $e\mathcal{R}f$.

Suppose the converse: that there exist two idempotents $e \neq f$ with $e\mathcal{R}f$. Then $\{e, f\}$ is a 2-element right zero semigroup. Restricting the action of $\overline{e}$ and $\overline{f}$ to $\{e, f\}^\infty$ yields the automaton $\mathcal{C}^*(\{e, f\})$. By Lemma 2.4, it follows now that $\langle \overline{e}, \overline{f}\rangle$ is a free semigroup of rank 2, a contradiction.

($\Leftarrow$). We will prove by induction on $|S|$ that if S is $\mathcal{H}$-trivial and every its $\mathcal{R}$-class contains at most one idempotent, then $\mathbf{C}^*(S)$ is finite. The base case $|S| = 1$ is obvious. Assume that we have proved this for all such semigroups of size $\leq n$. Take now any semigroup S with the assumption of sufficiency of the theorem such that $|S| = n + 1$.

Let M be the set of all maximal $\mathcal{D}$-classes in S, and let I be the complement of all these $\mathcal{D}$-classes in S. Suppose first that $I = \varnothing$. Then S consists of a single $\mathcal{D}$-class and so S is a Rees matrix semigroup $\mathcal{M}[G; X, Y; P]$ for some group G and a $Y \times X$-matrix $P = (p_{yx})_{Y\times X}$. Since S is $\mathcal{H}$-trivial, we have that $G = \{1\}$. Moreover, by [3, Theorem 3.4.2], we may assume that there exists a column in P consisting entirely of the element 1. Since columns in P correspond to $\mathcal{R}$-classes in S, we obtain that $|X| = 1$. So, again by [3, Theorem 3.4.2], it now holds that $S = Y$ is a left zero semigroup. Then by Lemma 2.3, $\mathbf{C}^*(S)$ is trivial.

So in the remainder of the proof we may assume that $I \neq \varnothing$. Obviously I is an ideal in S. We prove that $\mathbf{C}^*(S) = \langle \overline{S}\rangle$ is finite in the following four steps.

Step 1: $\langle \overline{\mathbf{I}}\rangle$ is finite.

It suffices to prove that there are finitely many products $\overline{i}\cdot\overline{i_1}\cdots\overline{i_k} \in \langle \overline{I}\rangle$ for any fixed $i \in I$. We have that $\overline{i}\cdot\overline{i_1}\cdots\overline{i_k}$ and $\overline{i}\cdot\overline{j_1}\cdots\overline{j_n}$ are distinct if and only if the restrictions of $\overline{i_1}\cdots\overline{i_k}$ and $\overline{j_1}\cdots\overline{j_n}$ on $S^\infty\overline{i}$ coincide. Notice that $S^\infty\overline{i} \subseteq I^\infty$. Obviously $\overline{i_1}\cdots\overline{i_k}$ and $\overline{j_1}\cdots\overline{j_n}$ act on I^∞ in the same way as the correspondent products from $\mathbf{C}^*(I)$ do. Now the claim of Step 1 follows from the induction hypothesis.

Step 2: $\overline{\mathbf{I}}\langle\overline{\mathbf{S}}\rangle^{\mathbf{1}} = \overline{\mathbf{I}} \cup \overline{\mathbf{I}}\langle\overline{\mathbf{S}}\rangle$ is finite.

Take a typical element $w = \overline{i}\cdot\overline{a_1}\cdots\overline{a_k} \in \overline{I}\langle\overline{S}\rangle$. Then for all $x \in S$, we have

$$q(\overline{i}\cdot\overline{a_1}\cdots\overline{a_k}, x) = \overline{ix}\cdot\overline{a_1xi}\cdot\overline{a_2xia_1}\cdots\overline{a_kxia_1\cdots a_{k-1}}.$$

Having that I is an ideal in S, we deduce that $q(\overline{i}\cdot\overline{a_1}\cdots\overline{a_k}, x) \in \langle\overline{I}\rangle$. Therefore, having that $\tau(w) = \rho_j$ for $j = i\cdot a_1\cdots a_k \in I$, we obtain $|\overline{I}\langle\overline{S}\rangle| \leq |I|\cdot|\langle\overline{I}\rangle|^{|S|}$. Step 2 now follows immediately.

Step 3: $\langle\overline{\mathbf{S}}\setminus\overline{\mathbf{I}}\rangle$ is finite.

Take $a_1, \ldots, a_k \in S \setminus I$. For all $x \in S$, we have

$$q_x = q(\overline{a_1}\cdots\overline{a_k}, x) = \overline{a_1x}\cdot\overline{a_2xa_1}\cdots\overline{a_kxa_1\cdots a_{k-1}}.$$

Obviously, to prove Step 3, it suffices to show that there are only finitely many expressions q_x for all $a_1, \ldots, a_k \in S \setminus I$ and $x \in S$. Moreover, by Step 2, it even suffices to show finiteness of the set P_0 of all such expressions q_x with additional requirement that $a_1x \in S \setminus I$ and $a_2xa_1 \in S \setminus I$. (Otherwise $a_2xa_1 \in I$ and so q_x comes from the finite set $\overline{S}\cdot\overline{I}\langle\overline{S}\rangle^1$.)

Observe that if $a_1, \dots, a_i, x$ do not come from the same $\mathcal{D}$-class from M, for some i, then $a_i x a_1 \cdots a_{k-1} \in I$. In particular, if $a_1 x$ and a_1 are not elements from the same $\mathcal{D}$-class in M, then $a_1 x \in I$ and so $q_x \in \overline{I}\langle\overline{S}\rangle^1$.

So, take $x \in S$ such that $a_1 x \in S \setminus I$ and $a_2 x a_1 \in S \setminus I$. Find the maximum number m such that $a_i x a_1 \cdots a_{i-1} \in S \setminus I$ for all $i \leq m$. Then, by the preceding paragraph, all $x, a_1, \dots, a_m$ are from the same $\mathcal{D}$-class in M. Now, $a_{m+1} x a_1 \cdots a_m \in I$ and so, by Step 2, to prove Step 3, it suffices to prove finiteness of the set $P_1 \subseteq P_0$:

$$P_1 = \{\overline{a_1 x} \cdot \overline{a_2 x a_1} \cdots \overline{a_k x a_1 \cdots a_{k-1}} \mid \ a_i x a_1 \cdots a_{i-1} \in S \setminus I \text{ for all } i \leq k\}.$$

Take a typical product $\overline{a_1 x} \cdot \overline{a_2 x a_1} \cdots \overline{a_k x a_1 \cdots a_{k-1}} \in P_1$. Then as we discussed above, $x, a_1, \dots, a_k \in D$ for some $\mathcal{D}$-class D from M. Moreover, $a_1 x, \dots, a_k x \in D$. Hence by Lemma 2.5 and Miller-Clifford Theorem, we have that each $\mathcal{H}$-class $L_{a_1} \cap R_x, \dots, L_{a_k} \cap R_x$ contains an idempotent. Then, by hypothesis, $a_1 \mathcal{L} \cdots \mathcal{L} a_k$.

Now, for all $i \leq k-1$, we have $a_i a_{i+1} \in D$ and so $L_{a_i} \cap R_{a_{i+1}}$ contains an idempotent. Having that $a_i \mathcal{L} a_{i+1}$ and by $\mathcal{H}$-triviality, we deduce that a_{i+1} is an idempotent. Then $a_1 = a_1 a_2 = \cdots = a_1 \cdots a_{k-1}$. Hence

$$q_x = \overline{a_1 x} \cdot \overline{a_2 x a_1} \cdots \overline{a_k x a_1 \cdots a_{k-1}} = \overline{a_1 x} \cdot \overline{a_2 x a_1} \cdots \overline{a_k x a_1}.$$

Finally, $a_1 x \in D$ and so $x a_1$, being the unique element of $L_{a_1} \cap R_x$, is an idempotent. Then for all $2 \leq i \leq k$, $a_i \mathcal{L} a_1 \mathcal{L} x a_1$, and since a_i is an idempotent, we obtain that $a_i x a_1 = a_i$. Therefore $q_x = \overline{a_1 x} \cdot \overline{a_2} \cdots \overline{a_k}$. Recall also that $a_2, \dots, a_k$ are $\mathcal{L}$-equivalent idempotents.

So that, to establish Step 3, it suffices to prove finiteness of the set

$$P_2 = \{\overline{a_1} \cdots \overline{a_k} \mid \ a_1, \dots, a_k \text{ are } \mathcal{L} - \text{equivalent idempotents from } S \setminus I\}.$$

Take a typical product $\overline{a_1} \cdots \overline{a_k} \in P_2$. Let $x \in S$ be arbitrary and let $q_x = q(\overline{a_1} \cdots \overline{a_k}, x)$. Then $q_x = \overline{a_1 x} \cdot \overline{a_2 x a_1} \cdots \overline{a_k x a_1}$. Consider now the following three cases:

Case 1: $(a_1 x, a_1) \notin \mathcal{D}$**.**

Then, as above, there are only at most $k_1 = |\overline{I}\langle\overline{S}\rangle^1|$ many such q_x-s.

Case 2: $a_1 x \mathcal{D} a_1$ **but** $(a_1 x, a_1) \notin \mathcal{L}$**.**

Let $y \in S$ be arbitrary. Then

$$q_{x,y} = q(q_x, y) = \overline{a_1 x y} \cdot \overline{a_2 x a_1 y a_1 x} \cdot \overline{a_3 x a_1 y a_1 x a_2 x a_1} \cdots .$$

We will prove that if $k > 2$, then $a_3 x a_1 y a_1 x a_2 x a_1 \in I$. Suppose the converse. Then $a_3 x a_1 y a_1 x a_2 x a_1 \mathcal{D} a_1$ and so $a_1 x, a_2, a_1 x a_2$ all lie in the same $\mathcal{D}$-class D_{a_1}. Then by Lemma 2.5, we obtain that $L_{a_1 x} \cap R_{a_2} = \{e\}$ contains an idempotent. But $a_1 \mathcal{L} a_2$ and $(a_1 x, a_1) \notin \mathcal{L}$, so that a_2 and e are $\mathcal{R}$-equivalent distinct idempotents, a contradiction.

Therefore $q_{x,y} \in \overline{S} \cup \overline{S}^2 \cup (\overline{S}^2 \cdot \overline{I}\langle\overline{S}\rangle^1)$. In turn, it implies that there at most k_2 elements q_x, where k_2 depends only on S.

Case 3: $a_1 x \mathcal{L} a_1$**.**

Then in particular we have $x \mathcal{L} a_1$ and x is an idempotent. Hence $a_1 x = a_1$, $a_2 x a_1 = a_2, \dots, a_k x a_1 = a_k$. Therefore $q_x = \overline{a_1} \cdots \overline{a_k}$.

So, from Case 3 it follows that $\overline{a_1} \cdots \overline{a_k}$ is uniquely determined by $\tau(\overline{a_1} \cdots \overline{a_k})$ and q_x-s with $x \in S$ such that $(a_1 x, a_1) \notin \mathcal{L}$. Thus from Cases 1 and 2 we have

$$|P_2| \leq |S| \cdot (k_1 + k_2)^{|S|}.$$

So, P_2 is finite and thus Step 3 is established.

Step 4: $\langle\overline{\mathbf{S}}\rangle$ is finite.

Follows from $\langle\overline{S}\rangle = \overline{I}\langle\overline{S}\rangle^1 \cup \langle\overline{S}\setminus\overline{I}\rangle \cup \langle\overline{S}\setminus\overline{I}\rangle\overline{I}\langle\overline{S}\rangle^1$, by Steps 2 and 3. $\square$

References

[1] A. J. Cain, V. Maltcev, Automaton semigroups, submitted.

[2] A. H. Clifford, G. B. Preston, The algebraic theory of semigroups. Vol. I, *Mathematical Surveys, No. 7 American Mathematical Society, Providence, R.I.* (1961).

[3] J. M. Howie, Fundamentals of semigroup theory, *Oxford Science Publications. The Clarendon Press, Oxford University Press, New York* (1995).

[4] A. Mintz, On the Cayley semigroup of a finite aperidoic semigroup, to appear in *Internat. J. Algebra and Comput.*.

[5] V. Nekrashevych, Self-similar groups, *Mathematical Surveys and Monographs* **117** *American Mathematical Society, Providence, RI* (2005).

[6] P. V. Silva, B. Steinberg, On a class of automata groups generalizing lamplighter groups, *Internat. J. Algebra Comput.* **15** (2005), no. 5-6, 1213–1234.